ORIGINAL ARTICLE

INTERNATIONAL JOURNAL OF ARTIFICIAL INTELLIGENCE

ISSN: 2692-515-x, Impact Factor: 9.23

American Academic publishers, volume 6, issue 01,2026

Journal: https://www.academicpublishers.org/journals/index.php/ijai

CAUSES AND CLINICAL FEATURES OF MENSTRUAL CYCLE DISORDERS IN ADOLESCENT GIRLS

Ismatova Farida Zokir kizi
Samarkand State Medical University
Intern's clinic

Abstract: Menstrual cycle disorders are among the most common gynecological problems encountered during adolescence. The period following menarche is characterized by physiological immaturity of the hypothalamic–pituitary–ovarian axis, which often results in irregular menstrual patterns. While many of these disturbances are transient and functional, some may indicate underlying endocrine, metabolic, or systemic conditions. Understanding the causes and clinical features of menstrual cycle disorders in adolescent girls is essential for early diagnosis, appropriate management, and prevention of long-term reproductive and psychological complications. This article reviews the major etiological factors and clinical manifestations of menstrual cycle disorders in adolescents based on modern medical knowledge.

Keywords: Adolescent health, menstrual disorders, puberty, hormonal imbalance, gynecology

Introduction

Adolescence represents a critical transitional period marked by rapid physical, hormonal, and psychological changes. The onset of menstruation is a key milestone of female pubertal development and reflects the maturation of complex neuroendocrine mechanisms. In the early years after menarche, menstrual irregularities are frequently observed and are often considered a normal part of development. However, persistent or severe menstrual disturbances should not be overlooked, as they may signal pathological conditions requiring medical attention. Menstrual cycle disorders can negatively affect quality of life, academic performance, and emotional well-being in adolescent girls, highlighting the importance of timely recognition and evaluation.

The most common cause of menstrual cycle disorders in adolescent girls is the immaturity of the hypothalamic–pituitary–ovarian axis. Inadequate coordination between gonadotropin-releasing hormone, luteinizing hormone, and follicle-stimulating hormone leads to anovulatory cycles, which manifest as irregular or abnormal uterine bleeding. Hormonal imbalance remains the central mechanism underlying most menstrual disorders during early adolescence.

Endocrine disorders such as polycystic ovary syndrome, thyroid dysfunction, and hyperprolactinemia play a significant role in the development of persistent menstrual irregularities. Polycystic ovary syndrome is particularly important due to its association with hyperandrogenism, insulin resistance, and long-term metabolic risks. Nutritional factors, including undernutrition, obesity, and eating disorders, also significantly influence menstrual regularity by altering hormonal secretion and energy balance.

Psychological stress and emotional disturbances are increasingly recognized as contributing factors. Academic pressure, family conflicts, and social stress can disrupt hypothalamic function, leading to delayed or irregular menstruation. Chronic systemic illnesses, excessive physical activity, and the use of certain medications may further interfere with normal menstrual regulation.

The clinical presentation of menstrual cycle disorders in adolescent girls is diverse and depends on the underlying cause. Common manifestations include irregular menstrual cycles,

ISSN: 2692-515-x, Impact Factor: 9.23
American Academic publishers, volume 6, issue 01,2026
Journal: https://www.academicpublishers.org/journals/index.php/ijai

prolonged intervals between periods, frequent bleeding episodes, or unpredictable menstruation. Some adolescents experience heavy menstrual bleeding, which may lead to fatigue, dizziness, and iron deficiency anemia.

Painful menstruation is another frequent complaint and may be primary in nature or associated with underlying pelvic pathology. In cases related to endocrine disorders, menstrual disturbances are often accompanied by additional signs such as acne, hirsutism, weight changes, or galactorrhea. Delayed menarche or secondary amenorrhea may indicate more serious systemic or hormonal conditions requiring thorough evaluation. The menstrual cycle in adolescent girls is regulated by a complex interaction between the hypothalamus, pituitary gland, and ovaries. During the early postmenarchal years, this regulatory system is still developing, which explains why menstrual irregularities are common in this age group. Ovulatory cycles may not be consistently established, leading to variations in cycle length, flow, and duration. In most cases, these changes are functional and gradually normalize as endocrine maturation progresses.

Hormonal imbalance is the central mechanism underlying most menstrual cycle disorders in adolescence. Inadequate pulsatile secretion of gonadotropin-releasing hormone results in insufficient stimulation of luteinizing hormone and follicle-stimulating hormone, causing anovulation. Anovulatory cycles often present with irregular bleeding patterns or prolonged menstrual intervals. Estrogen dominance without adequate progesterone support may lead to excessive endometrial proliferation and abnormal uterine bleeding.

Endocrine disorders represent an important pathological cause of persistent menstrual disturbances. Polycystic ovary syndrome is one of the most frequently diagnosed conditions in adolescent girls with menstrual irregularities. It is characterized by chronic anovulation, hyperandrogenism, and metabolic disturbances. Adolescents with this condition often present with infrequent menstruation, acne, excessive hair growth, and weight gain. Thyroid dysfunction, both hypothyroidism and hyperthyroidism, can also disrupt menstrual regularity by altering metabolic rate and hormonal feedback mechanisms.

Nutritional status plays a critical role in menstrual health during adolescence. Both undernutrition and obesity can negatively affect reproductive function. Insufficient caloric intake or eating disorders lead to decreased leptin levels and suppression of hypothalamic activity, resulting in delayed menarche or amenorrhea. Conversely, excess adipose tissue contributes to increased peripheral estrogen production, which may cause irregular or heavy menstrual bleeding. Rapid weight changes during adolescence further exacerbate hormonal instability.

Psychological and emotional factors are increasingly recognized as contributors to menstrual cycle disorders. Chronic stress activates the hypothalamic–pituitary–adrenal axis, increasing cortisol levels and inhibiting normal gonadotropin secretion. Academic pressure, family conflicts, anxiety, and depression are common stressors during adolescence and may manifest clinically as delayed, absent, or irregular menstruation. These functional disorders often coexist with somatic complaints such as headaches, fatigue, and sleep disturbances.

Clinical manifestations of menstrual cycle disorders vary widely. Some adolescent girls experience oligomenorrhea with prolonged intervals between menstrual periods, while others report frequent or unpredictable bleeding. Heavy menstrual bleeding is particularly concerning, as it may lead to iron deficiency anemia, reduced physical endurance, and impaired cognitive performance. Dysmenorrhea is another prevalent symptom and may significantly interfere with daily activities and school attendance.

ORIGINAL ARTICLE
INTERNATIONAL JOURNAL OF ARTIFICIAL INTELLIGENCE
ISSN: 2692-515-x, Impact Factor: 9.23
American Academic publishers, volume 6, issue 01,2026
Journal: https://www.academicpublishers.org/journals/index.php/ijai

In certain cases, menstrual disorders may be the first visible sign of underlying systemic disease. Chronic illnesses, excessive physical activity, and the use of medications such as antipsychotics or hormonal agents can disrupt normal menstrual patterns. Therefore, a comprehensive clinical evaluation that includes medical history, physical examination, and appropriate laboratory investigations is essential for accurate diagnosis.

Overall, menstrual cycle disorders in adolescent girls reflect a dynamic interaction between physiological maturation and external influences. Early recognition of abnormal patterns and identification of contributing factors are crucial for ensuring normal reproductive development and protecting long-term health.

The regulation of the menstrual cycle in adolescent girls depends on the gradual maturation of neuroendocrine pathways that connect the central nervous system with the reproductive organs. During puberty, the hypothalamus begins to secrete gonadotropin-releasing hormone in a pulsatile manner, which stimulates the pituitary gland to release luteinizing hormone and follicle-stimulating hormone. These hormones act on the ovaries to promote follicular development and ovulation. In adolescents, this coordination is often incomplete, resulting in irregular, anovulatory menstrual cycles that may persist for several years after menarche.

One of the most important mechanisms contributing to menstrual cycle disorders is estrogen–progesterone imbalance. In the absence of ovulation, progesterone production is insufficient, leading to prolonged estrogen exposure of the endometrium. This hormonal environment causes unstable endometrial growth and irregular shedding, clinically presenting as abnormal uterine bleeding. Such bleeding patterns are common in early adolescence but may become pathological if prolonged or excessive.

Endocrine pathologies significantly influence menstrual regularity in adolescent girls. Polycystic ovary syndrome is increasingly diagnosed in this age group due to changes in lifestyle and rising obesity rates. The syndrome involves altered ovarian steroidogenesis, insulin resistance, and elevated androgen levels, which interfere with follicular maturation. Menstrual irregularities associated with this condition are often chronic and may persist into adulthood if left untreated. Thyroid hormones also play a crucial role in menstrual regulation by influencing metabolic processes and reproductive hormone feedback. Even mild thyroid dysfunction can lead to cycle irregularities, emphasizing the need for hormonal screening in persistent cases.

Metabolic and nutritional factors exert a profound impact on menstrual health. Adequate energy availability is essential for normal hypothalamic function. Adolescents with low body fat percentage or restrictive eating behaviors often experience hypothalamic suppression, resulting in amenorrhea or delayed menstrual cycles. On the other hand, excess body weight alters insulin sensitivity and increases estrogen production from adipose tissue, contributing to irregular and heavy menstrual bleeding. These metabolic disturbances are often accompanied by long-term risks such as infertility and cardiovascular disease.

Psychosocial influences are particularly significant during adolescence. Emotional stress affects the hypothalamic–pituitary–ovarian axis through increased secretion of stress hormones, which inhibit reproductive hormone release. Adolescents facing chronic stress may present with functional menstrual disorders without identifiable organic pathology. These disturbances highlight the close relationship between mental health and reproductive function, underscoring the importance of a holistic clinical approach.

The clinical spectrum of menstrual cycle disorders in adolescent girls includes amenorrhea, oligomenorrhea, polymenorrhea, and menorrhagia. Heavy menstrual bleeding is a major clinical concern due to its association with iron deficiency anemia, weakness, impaired concentration, and reduced academic performance. Painful menstruation is also common and

ISSN: 2692-515-x, Impact Factor: 9.23
American Academic publishers, volume 6, issue 01,2026
Journal: https://www.academicpublishers.org/journals/index.php/ijai

may be exacerbated by increased prostaglandin production during anovulatory cycles. Severe dysmenorrhea warrants evaluation to exclude secondary causes such as congenital anomalies or inflammatory conditions.

From a diagnostic perspective, careful assessment of menstrual history is essential. Evaluation should focus on age at menarche, cycle regularity, duration and volume of bleeding, associated symptoms, and lifestyle factors. Physical examination and laboratory investigations help differentiate functional disorders from pathological conditions. Early diagnosis allows for targeted interventions, including lifestyle modification, hormonal therapy, and psychological support.

In modern clinical practice, management of menstrual cycle disorders in adolescents emphasizes individualized care. Education and reassurance are often sufficient for functional disturbances, while persistent or severe disorders require multidisciplinary management. Addressing menstrual health during adolescence plays a critical role in ensuring normal reproductive development, preventing chronic gynecological disorders, and improving overall quality of life.

Conclusion

Menstrual cycle disorders in adolescent girls are common and multifactorial in origin. While many disturbances reflect normal physiological maturation, persistent or severe abnormalities warrant careful assessment to exclude underlying pathology. Early identification and appropriate management are crucial to prevent long-term reproductive, metabolic, and psychological consequences. A comprehensive approach that considers hormonal, nutritional, psychological, and lifestyle factors is essential in promoting optimal reproductive health during adolescence.

References

1. Diaz A, Laufer M. Menstrual disorders in the adolescent age group. Pediatric Clinics of North America. 2019;66:55–70.
2. Hickey M, Balen A. Menstrual disorders in adolescence: investigation and management. Human Reproduction Update. 2020;26:356–371.
3. Klein D, Poth M. Amenorrhea in adolescents: a diagnostic approach. American Family Physician. 2018;98:39–48.
4. Ibáñez L, Oberfield S, Witchel S. Anovulation in adolescent girls. Endocrine Reviews. 2020;41:671–705.
5. Deligeoroglou E, Karountzos V. Menstrual disturbances during puberty. Annals of the New York Academy of Sciences. 2019;1457:52–58.
6. Apter D. Development of the hypothalamic–pituitary–ovarian axis. Annals of the New York Academy of Sciences. 2018;1135:9–21.
7. Rosenfield R. Clinical review: adolescent anovulation. Journal of Clinical Endocrinology and Metabolism. 2021;106:1214–1226.