



## EFFECT OF MICRONUTRIENTS IN THE BACKGROUND OF MINERAL FERTILIZERS ON FORMATION OF YIELD ELEMENTS OF SOYBEAN VARIETIES

*Khayrullaev Sardor Shamsiddin ugli<sup>1</sup>, Usmonova Shokhista Usmon kizi<sup>2</sup>*

<sup>1</sup>Assistant teacher of Tashkent state agrarian university, Orcid ID: <https://orcid.org/0000-0001-5670-6449>

<sup>2</sup>Researcher of Rice Research Institute, Orcid ID: <https://orcid.org/0009-0005-3401-3031>

**ABSTRACT:** In this paper, Information is given about effect of micronutrients in the background of mineral fertilizers on formation of yield elements of soybean varieties. According to the information, under the influence of micronutrients, it was determined that it was observed an increase in the quantitative indicators of all yield elements and high results in the formation of yield elements were recorded when sulfur and manganese micronutrients were used in medium norms compared to other norms

**Keywords:** background, formation of yield elements, development, growth, macro and micronutrients, manganese, soybean, sulfur, varieties.

### INTRODUCTION

Soybean production is a branch which appeared in an ancient period. According to sources, it appeared in the northern, central and southern regions of China 5000 years ago. The development of soy farming is closely related to the development of agriculture. Soybean is a very ancient crop. Studying the diversity of soybean species and forms, scientists believe that it was formed mainly in three centers: Southeast Asia, Australia and East Africa. But many scientists recognize the center of China as the origin of the plant - China, Korea, India, Japan. Soya appeared in Europe and the USA in 1712. For many centuries, soybean and its products served as the main source of protein for the people of eastern countries. Soybean was cultivated in China 6-7 thousand years ago. It has been cultivated since ancient times in Japan, India and Indonesia, Vietnam and other Asian countries. Soybean entered the Far East from China 3 thousand years ago. Soybeans have long been grown in Russia's Amur region, Khabarovsk and Primorsky regions, where natural and climatic conditions are favorable. Later, it began to be cultivated in Georgia, Ukraine, the Kuban and the North Caucasus. During evolution, soybean species and varieties were grown and adapted to different soil and climate conditions far from their homeland, the morphological and biological characteristics of the plant partially changed [1].

Soybean plant plays an important role in solving some of the existing problems in agriculture. Oil, margarine, cheese, milk, flour, confectionery products are obtained from soybeans. Soybean flour is added to bread and sausage products and increases the nutritional value, taste and strength of these products. Soybean products are recommended for diabetes. Soybeans are rich in minerals such as potassium, calcium, and phosphorus [2].

Kh.N. Atabaeva., F.B. Namozov., A.A. Kurbanov and S.Sh. Khayrullaev in their experiments conducted in 2018-2020, when they applied micronutrients to the soybean crop, micronutrients affected the height of the soybean stem, leaf, root development, nodule formation, grain quality and productivity, and provided a high yield [3].

According to R. Jo'raeva., J. Toshpol'atov., A. Iminov., Kh. Bozorov and L. Zaynitdinova, S. Khatamov and

S. Sh. Khayrullaev, in their experiments conducted in 2015-2017, soybean plant mineral fertilizers and belonging to the rhizobium group it was observed that the yield increased by 12.6-12.8 c/ha when exposed to azotobacteria strains compared to the control variant [4, 7].

According to Khayrullayev Sardor Shamsiddin ugli (2021), the application of micronutrients in the suspension method 2 times during the application period of soybean varieties in the conditions of meadow-swamp soils provides an increase in grain quality [9]. According to data of Atabayeva Khalima Nazarovna, Khayrullaev Sardor Shamsiddin o'g'li, and Usmonova Shohista Usmon qizi (2020), sulfur has a positive effect on the branching of soybean varieties on the background of mineral fertilizers, and in 2018 the number of branches in the variety "Orzu" increased by 0.8-1.3 compared to the control option due to the micro element sulfur. In the "Nafis" variety, this figure was 0.3-0.4, and good results were obtained from medium and high sulfur norms. In 2019, these indicators increased by 0.3-0.7 in the variants of sulfur compared to the control in the "Orzu" variety, increased by 0.1-0.3 in the "Nafis" variety, and good results were obtained from the medium and high norms of sulfur [6]. According to Iminov Abduvali Abdumannobovich, Khayrullayev Sardor Shamsiddin ugli, et al, Nitragine treatment of soybean and mung bean seeds before sowing had a positive effect on seed germination under both laboratory and field conditions, the germination rate of seeds in the laboratory under the conditions of cotton cultivation in the following year under the background of non-treatment by nitragine before sowing the seeds of soybean and mungbean crops grown as a secondary crop after winter wheat was 0.3-1.3%, and field fertility was 0.2-0.8% higher. Also, it was found that the use of phosphorus and potassium fertilizers in soybean and mung bean crops grown as a secondary crop was 0.6-1.0% higher in the laboratory, and 0.6-0.7% higher in the field than in the control options without mineral fertilizers in studies [5]. According to Umarova Nigora Sadridinovna, Bo'riboyev Bekzod Yetmish ugli, Khayrullayev Sardor Shamsiddin ugli, Usmonova Shokhista Usmon kizi, & Turdaliyeva Shokhista Tulkinjon kizi, the demand of the soybean plant for mineral fertilizers, it was observed that when NPK and liquid fertilizer were used together, all the biometric parameters and yields of the plant increased by varieties compared to other methods. The use of mineral fertilizers in different ways in typical sierozem soil conditions affects the grain yield of local and foreign varieties. In other words, the average yield of medium-ripe soybean varieties "Nafis" was 43.4 c / ha, "Vilana" was 42.4 c / ha, and the best way to increase the yield is to apply fertilizers as NPK in combination with liquid fertilizer [10]. According to data of Khayrullayev Sardor Shamsiddin o'g'li and Usmonova Shohista Usmon qizi, the location of the lower first pod in soybean varieties is 12.8-15.9 cm in Orzu variety, 3-3.1 cm in Radimax stimulator, 2.2-2.4 cm in Gummat stimulator, 2.1 cm in Tecamin stimulator and 3.1 cm in Algora stimulator was found to be high. The most effective results were observed in Radimax, Gummat and Algora bio-simulators, and the location of the lower first pod was detected 14.7-17.6 cm in the "Nafis" variety, which was 2.5-2.9 cm higher in the Radimax stimulator, 2.2-2.5 cm higher in the Gummat stimulator, 2.1 cm higher in the Tecamine stimulator, and 2.4 cm higher in the Algora stimulator than in the control variant. The most effective results were observed in Radimax, Gummat and Algora biosimulators [8]. According to Atabayeva, K. N., Umarova, N. S., Yakubov, S., & Khayrullaev, S. S, positive results were obtained from moderate levels of sulphur and manganese, and low levels of iron. Macro and micronutrients had a positive effect on soy yield. An additional 7.6 quintals (q)/ha was harvested in exchange for macro fertilizer. Compared to the background variant, the yield was 4.6-8.3 q/ha for sulphur and 4.9-9.8 q/ha for manganese. The yield of the iron element was lower than that of the background variant. Grain quality has changed in exchange for macro and micronutrients. In exchange for mineral fertilizers, this figure increased by 2.4%. In exchange for the element sulphur, the protein increased by 3.1-5.8%; an increase of 4.4-8.4% was observed in exchange for the element manganese. It was noted that the protein increased by 7.9-8.7% in exchange for the element iron [11]. Khayrullayev, S. S., & Kizi Usmonova, S. U explained that mineral fertilizers and sulfur microelements activate the symbiotic activity of soybean variety "Orzu", averaging 32.4-42.3 million nodules per hectare, the number of nodules due to the background of mineral fertilizers increased by 13.6%, and there was an increase of 19.4-23.4% due to sulfur, as well as an average weight of nodules was 6.46-9.56 c / ha, the weight of nodules increased by 5.3% due to mineral fertilizers, and 17.1-32.4% due to sulfur. During the application period, 6.46-9.56 c / ha of nodules mass was accumulated per hectare according to the studied variants, which contributes to the increase of nitrogen and organic matter in the soil and a slight increase in biological efficiency [12]. Usmonova Sh.U, Khayrullaev S.Sh, Shomuqimov

N.N., & Gaynanova A.F, said that the influence of stimulants on soybeans affected the weight of 1000 grains of Vilana cultivar, under the influence of Gummat stimulator this figure was 2.2-7.4 grams higher than on basis of mineral fertilizers (Background), and under the influence of Rival stimulator-3.0-6.0 grams [13]. According to Khayrullaev S. S, In the variant, where not used mineral fertilizers and micronutrients, the leaf area in the control variant of the Orzu variety of soybean was 51.1 thousand m<sup>2</sup> / ha. Under the influence of microelements, the leaf area of Orzu was 59.1-64.6 thousand m<sup>2</sup> / ha. The highest rates of exposure to micronutrients were observed with medium use of sulfur and manganese. Under influence macro and micro fertilizers, the leaf area of Orzu variety increased from 4.0 to 13.5 thousand m<sup>2</sup> / ha, or from 7.3 to 20.9% [14].

It was found that when the plants were exposed to urea at different rates, they had an effect on their biometric indicators, and when the urea rate was increased to 15 grams, all biometric indicators were better than other options (Umarova Nigora Sadridinovna, et.al. 2023) [15]. The application of stimulants to plants increases their resistance to abiotic factors. All the stimulants in the experiment showed their effect for the preservation of the plant, and the foliar feeding variant of the fulvogummat stimulator showed a better result than the other stimulants (Usmonova Shokhista Usman kizi, et.al. 2023) [16].

Usmonova Shokhista and other scientists noticed that the growth process has a great role in the formation of the plant's fruit, and the use of stimulants in the experiment showed a positive effect on the growth process of the plant. The studied stimulants have an effect on the growth process, among the stimulants Fulvogummat stimulator was 164.9 cm tall when fed from the leaf, and recorded a good indicator among the variants (Usmonova Shokhista Usman kizi and et.al. 2023) [17].

## METHODS AND MATERIALS

Experiments are carried out in field and laboratory conditions. In the research "Methods of conducting field experiments" (T.UzPITI 2007), "Methodology of field experiment (B.Dospekhov, 1985), "Methodology of the State variety testing of agricultural crops" (1985, 1989), "Methods of agrochemical, agrophysical studies of the soil of Central Asia" (1988) methods are used.

### Varieties used in the experiment

**Orzu the variety** is an early-ripening variety, 35-40 days pass from planting to flowering, 95-110 days to the ripening period, the stem branches. The stem grows upright, the bush is porous, the height of the stem can be up to 50-70 cm. The leaf is three-lobed, large, light green. The foliage of the plant is average, the leaves are symmetrical. The length of the leaf band is 10 cm. 75% of the leaves are shed when fully ripe. The flower is white, there are 2-7 flowers in the inflorescence. The pods are gray, small, from 2.4 cm long up to 4.0 cm. Pods do not split when ripe, on average about 40 pods are formed per bush. The grain is average, the weight of 1000 grains is 120-130 g. Grain yield in irrigated lands is 32 centners per hectare. when planted as repetitive plant 10-20 c grain yield is taken. Grain contains 25% oil and 36-38 % protein. Authors: Rakhmanov A.R, Yunusov B.K, Tulaganov N., Burigina O.V.

**Nafis variety.** The variety was created by the method of individual selection at the Rice Research Institute of Uzbekistan.

The growing period is 115-120 days. The height of the plant is 145-150 cm. The location of the lower pod is 14-16 cm, the number of branches is 2-4, the number of pods in one plant is 120-130, the number of grains in one pod is 2-4.

Grain quality and technological parameters: weight of 1000 seeds is 165-175 g. The protein content of grain is 40-41%, the oil content is 25-27%. Resistant to lodging, shedding, diseases and mechanized harvesting.

Yield: 30-32 c/ha grain yield and 250-300 c/ha blue mass can be obtained from the variety under favorable conditions.

### Place of experiment, conditions and agrotechnical measures

The experiments were conducted in the scientific experimental fields of the Rice Research Institute in Tashkent region.

The soil layers are swamp type soils characteristic of an oasis. There are also large and small stones and sand mixtures in different depth layers. These soils derive from the typical excess moisture conditions of the

left bank of the river and are ideal for rice cultivation. The soil is grassland. The soil of the experimental field is not saline, the tiller layer is 30- 40 cm. The pH of the solutions in the soil is 6.8-7.3 pieces, and it is heavy clay according to its mechanical composition. Experiments are being conducted in 4 checks of 12 maps.

Prior to planting, the background was established in the program, in which 50 kg of nitrogen, 100 kg of phosphorus and 70 kg of potassium were applied. Planting method is wide rows, row spacing is 70 cm, bush spacing is 5 cm. Nitragin was not used because soybeans are always grown at the Rice Institute and the soil contains Rhizobium bacteria.

The experimental field was irrigated 2 times during the period of growth. Cultivation was carried out 2 times in the experimental field with the help of equipment. Soybean varieties were fed in 3 different ways, suspension was used.

## RESULTS AND DISCUSSION

One of the main factors determining the productivity of leguminous crops is the degree of formation of their yield elements (the number of pods and grains) and the nature of their development.

Before harvesting, the composition of the crop is determined. In the field experiments, when the soybean crop was ripe, we analyzed the biometric measurements (stem height, number of branches, total stem weight, number and weight of pods, number and weight of grains) and it was found that there are differences between the varieties (see table 1).

Based on the 3-year average data of our experience, the stem height of the Orzu variety was 109.0-115.1 cm among the variants. The number of branches was 1.36-2.50 pieces among the options. The number of pods is one of the main elements of the yield. The number of pods per plant was 48.9 pieces in the control option, and 4.5 pieces more than the control option in the background option with mineral fertilizers. In the options where sulfur micronutrient was applied against the background of mineral fertilizers, the number of pods was 12.9, 19.0 and 17.0 more than the control option in accordance with the norms.

**Table 1**

Application norms of micronutrients, g/l	Stem height, cm	Number of branches, piece	Number of pods in one bush plant, piece	Number of grain in one bush plant, g	Weight of pods in one bush plant, piece	Weight of grain in one bush plant, g	Weight of pods in countable rows, g	Weight of grain in countable rows, g	Weight of stem in countable rows, g
<b>Orzu variety</b>									
Without fertilizers-(control)	109.0	1.36	48.9	122.3	23.1	17.9	2755.0	1313.7	5833.6
Background-N <sub>50</sub> P <sub>100</sub> K <sub>70</sub> kg/ha	110.4	1.53	53.4	123.5	24.3	19.2	2852.7	1845.7	5941.5
Background+S <sub>90</sub>	110.5	1.93	61.8	124.1	27.7	19.7	3043.8	2170.0	6787.9
Background+S <sub>180</sub>	113.8	2.50	67.9	146.2	29.2	20.8	3307.5	2496.7	7039.1
Background+S <sub>270</sub>	113.2	2.23	65.9	130.2	28.2	19.8	3262.7	2291.3	6753.0
Background+Mn <sub>150</sub>	113.0	1.66	59.1	135.4	30.4	19.3	3108.0	2188.7	7173.1
Background+Mn <sub>300</sub>	115.1	2.00	73.9	178.2	39.0	26.6	3933.8	2464.0	9669.4
Background+Mn <sub>450</sub>	113.5	1.83	64.0	149.8	33.3	22.5	3412.4	2242.3	9513.1
<b>Nafis variety</b>									



Without fertilizers-(control)	149.1	0.82	60.8	174.5	44.2	31.3	1945.0	1521.3	5320.9
Background-N <sub>50</sub> P <sub>100</sub> K <sub>70</sub> kg/ha	151.4	1.02	66.8	177.7	46.2	32.9	2354.4	2023.0	5499.4
Background+S <sub>90</sub>	152.9	1.35	70.3	186.1	47.8	34.4	3007.1	2454.7	6027.4
Background+S <sub>180</sub>	158.0	1.95	85.7	205.6	53.6	37.3	3706.7	2578.3	7431.2
Background+S <sub>270</sub>	153.8	1.48	77.7	202.2	51.7	36.6	3550.9	2496.7	6535.8
Background+Mn <sub>150</sub>	157.0	1.05	72.2	196.2	48.0	34.0	3542.8	2454.7	6339.3
Background+Mn <sub>300</sub>	160.3	1.46	75.9	198.4	50.5	36.2	3894.1	2695.0	8272.4
Background+Mn <sub>450</sub>	158.7	1.20	72.5	197.0	49.9	35.4	3673.1	2576.0	6501.5

### **The effect of micronutrients in the background of mineral fertilizers on the formation of yield elements of soybean varieties, (2018-2020)**

In the options where manganese micronutrient was used in the background of mineral fertilizers, the number of pods was increased by 10.2, 25.0 and 15.1 pieces more than the control option in accordance with the norms.

The number of grains per plant varied from 122.3 to 178.2 among the variants. A high result of 146.2 and 178.2 pieces was recorded in the medium norms of sulfur and manganese micronutrients.

Pod weight per plant varied from 23.1 g to 39.0 g among variants. It was observed that the weight of grain per plant varied from 17.9 g to 26.6 g among the variants. It was noted that the weight of beans in the calculated row changed from 2755.0 grams to 3933.8 grams.

Grain weight in the calculated row is one of the important elements of the harvest. The grain weight in the calculated row was 1313.7 g in the control variant. The background option with mineral fertilizers was 532 g more than the control option. In the variants using sulfur micronutrient, the grain weight in the calculated row is 856.3 more than the control variant in accordance with the norms; It increased by 1183.0 and 977.6 g. The weight of the grain in the calculated row in the options using manganese micronutrient is 875.0 more than the control option in accordance with the norms; It increased by 1150.3 and 928.6 g.

The stem weight of plants means additional yield. Stem weight varied from 5833.6 grams to 9669.4 grams. The stem weight was 5833.6 g in the control variant. A good indicator was noted in the medium norm rate of sulfur micronutrient, which was 1205.5 g higher than the control option. A good indicator of manganese micronutrient was also noted in the medium, and it was 3835.8 g higher than the control option.

Based on the 3-year average data of our experience, the stem height of the "Nafis" variety was 149.1-160.3 cm among the variants. The number of branches was 0.82-1.95 among the options.

The number of pods is one of the main elements of the crop. The number of pods per plant was 60.8 pieces in the control option, and 6.0 pieces more than the control option in the background option with mineral fertilizers. In the variants using sulfur micronutrient, the number of pods is 70.3 in accordance with the norms; It was 85.7 and 77.7 pieces. In the variants with manganese micronutrient, the number of pods is 72.2 75.9 and 72.5 pieces in accordance with the norms.

The number of grains per plant varied from 174.5 to 205.6 among the variants. Sulfur micro-fertilizers and manganese micronutrients showed a high result with 205.6 and 198.4 pieces.

The pod weight per plant varied from 44.2 g to 53.6 g among the variants. The grain weight per plant varied from 31.3 to 37.3 g among the variants. The weight of beans in the calculated row varied from 1945.0 grams to 3706.7 grams.

Grain weight in the calculated row is one of the important elements of the harvest. The grain weight in the calculated row was 1521.3 g in the control variant. The background option with mineral fertilizers was 501.7 g more than the control option. In the variants with sulfur micronutrient, the grain weight in the calculated row is 933.4 1057 and 975.4 g more than the control variant in accordance with the norms. The weight of the grain in the calculated row in the options with manganese micronutrient is increased by 933.4, 1173.7 and 1054.7 g more than the control option in accordance with the norms.

Stem weight varied from 5320.9 grams to 8272.4 grams depending on the options. The stem weight was 5320.9 g in the control variant. A good indicator was noted in the medium norm standard of sulfur micronutrient, which was 2110.3 g higher than the control option. The same pattern was observed in the

manganese micronutrient, and a good indicator was noted in the medium norm, that is, it was 2951.5 g higher than the control option (see Table 1).

## CONCLUSION

Thus, under the influence of micronutrients, it was determined that it was observed an increase in the quantitative indicators of all yield elements and high results in the formation of yield elements were recorded when sulfur and manganese micronutrients were used in medium norms compared to other norms.

## REFERENCES:

1. Atabaeva H.N., Umarova N.S. – “Soybean biology”. - textbook, “Navruz” publishing house, 2020, 14.5 p.t., B.5-8, P.11-12. (In Uzbek language)
2. Atabaeva Kh.N. - Soybean. Monography // National encyclopedia of Uzbekistan . state scientific publishing house. 2004. P.96. (In Uzbek language)
3. Nazarovna, AK, Bakhromovich, NF, Alavkhonovich KA, Ugli KSS. Effects of Sulfur and Manganese Micronutrients on the Yield of Soybean Varieties. *Agricultural Sciences*, 2020;11: 1048-1059. <https://doi.org/10.4236/as.2020.1111068>
4. Juraeva R., Tashpulatov J., Iminov A., Bozorov X., Zaynitdinova L., & Kukanova S. (2020). Efficiency Of Symbiotic Nitrogen Fixation Of Soy Nodule Bacteria After Preservation. *Plant Cell Biotechnology And Molecular Biology*, 21(61-62), 7279. Retrieved from <https://www.ikprress.org/index.php/PCBMB/article/view/5644>.
5. Abdumannobovich, I. A., Ugli, K. S. S., Ergashboyevna, K. D., Ugli, H. S. R., & Tursinalievich, K. S. (2021). Effects Of Nitragine And Mineral Fertilizer Rates Used On Repeated Legume Crops On Seed Germinability. *Plant Cell Biotechnology And Molecular Biology*, 22(35-36), 251-258. Retrieved from <https://www.ikprress.org/index.php/PCBMB/article/view/6342>.
6. Atabayeva Khalima Nazarovna1, Khayrullaev Sardor Shamsiddin o’g’li, and Usmonova Shohista Usmon qizi, “The Effect Of Sulphur On The Branching Of Soybean Varieties”, *STJITT*, vol. 1, no. 2, pp. 4-6, Aug. 2020.
7. Iminov, A. A., Hatamov, S. R. O., & Khayrullaev, S. S. O. (2020). Effect Of Nitragine And Mineral Fertilizers On Soil Microbiological Properties In Planted As Secondary Legume Crops. *The American Journal of Agriculture and Biomedical Engineering*, 2(08), 169-172. <https://doi.org/10.37547/tajabe/Volume02Issue08-22>
8. Khayrullayev Sardor Shamsiddin o’g’li and Usmonova Shhista Usmon qizi, “The Effect Of Stimulators On The Location Of The First Pod Of Soybean Varieties”, *STJITT*, vol. 1, no. 2, pp. 7-9, Aug. 2020.
9. Khayrullayev Sardor Shamsiddinugli. Effect Of Sulfur And Manganese Micro Nutrients On Grain Quality Of Soybean Varieties. *ACADEMICIA: An International Multidisciplinary Research Journal* <https://saarj.com>. Vol. 11, Issue 3, March 2021. DOI: 10.5958/2249-7137.2021.00582.6. P.63-68.
10. Umarova Nigora Sadrididinovna, Bo’riboyev Bekzod Yetmish ugli, Khayrullayev Sardor Shamsiddin ugli, Usmonova Shokhista Usmon kizi, & Turdaliyeva Shokhista Tulkinjon kizi. (2021). Effect Of Mineral Fertilizers On Morphophysiological Characteristics Of Local And Foreign Soybean Varieties. *European Journal of Agricultural and Rural Education*, 2(4), 39-42. Retrieved from <https://scholarzest.com/index.php/ejare/article/view/611>.
11. Atabayeva, K. N., Umarova, N. S., Yakubov, S., & Khayrullaev, S. S. (2021, December). Influence of trace elements on soybean yield in grassland-swamp soils. In *IOP Conference Series: Earth and Environmental Science* (Vol. 939, No. 1, p. 012049). IOP Publishing.
12. Ugli Khayrullayev, S. S., & Kizi Usmonova, S. U. (2021). Effect Of Sulfur Micronutrient On Symbiotic Activity Of “Orzu” Variety Of Soybean. *The American Journal of Agriculture and Biomedical Engineering*, 3(05), 49–59. <https://doi.org/10.37547/tajabe/Volume03Issue05-10>
13. Usmonova Sh.U, Khayrullaev S.Sh, Shomuqimov N.N, & Gaynanova A.F. (2022). Effect Of Stimulators On Thousand Piece Grain Weight Of “Vilana” Cultivar Of Soybean. *Central Asian Journal of Theoretical and Applied Science*, 3(4), 153-157. Retrieved from <https://cajotas.centralasianstudies.org/index.php/CAJOTAS/article/view/467>
14. Khayrullaev S. S. Influence of sufur and manganese micronutrients on formation of leaf area of “Orzu”

cultivar of soybean //ISJ Theoretical & Applied Science, 03 (107). – 2022. – C. 287-191.

15. Umarova Nigora Sadridinovna, & Khayrullaev Sardor Shamsiddin ugli. (2023). The Effect Of Using Different Urea Norms In Suspension Method On Biometric Indicators Of Soybean. Spectrum Journal of Innovation, Reforms and Development, 15, 159–164. Retrieved from <https://sjird.journalspark.org/index.php/sjird/article/view/665>

16. Usmonova Shokhista Usman kizi, Atabaeva Halima Nazarovna, & Khayrullaev Sardor Shamsiddin ugli3. (2023). The Effect Of Stimulants On The Number Of Bushes And Preservation Level Of The Nafis Variety Of Soybean. Spectrum Journal of Innovation, Reforms and Development, 15, 177–184. Retrieved from <https://sjird.journalspark.org/index.php/sjird/article/view/668>

17. Usmonova Shokhista Usman kizi, Atabaeva Halima Nazarovna, & Khayrullaev Sardor Shamsiddin ugli. (2023). The Effect of Stimulants on the Stem Height of the Nafis Variety of Soybean. Texas Journal of Multidisciplinary Studies, 20, 86–91. Retrieved from <https://zienjournals.com/index.php/tjm/article/view/3988>