



PATHOGENETIC BASIS OF TREATMENT IN RECURRENT VARICOCELE

Bobokulov Nurillo Asatovich

Samarkand State Medical University

Nazarov Murodjon Normamatovich

Samarkand State Medical University

Vohidov Sanjar Qudratullayevich

Samarkand State Medical University

Abstract: Recurrent varicocele is a common urological disorder that negatively affects male reproductive function and fertility. It develops after previous surgical or minimally invasive interventions, resulting in persistent venous reflux, testicular hypoxia, and oxidative stress, which collectively impair spermatogenesis. Accurate diagnosis through clinical examination, Doppler ultrasonography, and advanced imaging modalities is essential for identifying recurrence and guiding treatment. Management strategies should be based on pathogenetic principles, incorporating microsurgical, endovascular, and conservative approaches tailored to patient anatomy, prior interventions, and fertility goals. Surgical precision, prevention of venous stasis, and postoperative monitoring are crucial for reducing recurrence rates. Adjunctive pharmacological therapies, including antioxidants and lifestyle modifications, support testicular function and enhance reproductive outcomes. This review highlights the pathophysiological mechanisms underlying recurrence, evaluates current treatment options, and emphasizes evidence-based strategies to preserve testicular function and improve quality of life in affected men.

Keywords: Recurrent varicocele, Pathogenesis, Venous reflux, Testicular hypoxia, Oxidative stress, Microsurgical treatment, Endovascular therapy, Male fertility.

Introduction

Varicocele is a pathological dilation of the testicular veins associated with venous reflux, representing one of the most common disorders of the male reproductive system. Epidemiological studies indicate that varicocele affects approximately 15–20% of the male population and can negatively impact spermatogenesis due to impaired venous blood flow. Recurrent varicocele refers to the reappearance of clinical symptoms following previous surgical or minimally invasive treatment, presenting both clinical and psychological challenges for patients. The main factors contributing to varicocele recurrence include deficiencies in the venous valve system, pathological structures of collateral veins, limitations of previously performed surgical techniques, and, in some cases, anatomical variations. From a pathogenetic perspective, recurrent varicocele is characterized by disrupted blood circulation and prolonged venous stasis, leading to testicular hypoxia, oxidative stress, and impaired function of testicular cells. Furthermore, preserving reproductive potential and testicular function in patients with recurrent varicocele requires modern therapeutic approaches based on anatomical and pathophysiological mechanisms, including surgical, endovascular, and conservative methods. Prevention and early detection of recurrence are also closely linked to pathogenetically grounded comprehensive interventions. This article analyzes the mechanisms of recurrent varicocele



development, its clinical significance, and therapeutic strategies based on pathogenetic principles, providing evidence-based recommendations for management.

Relevance

Varicocele is a common urological disorder among men that can lead to impaired reproductive function, reduced sperm quality, and in some cases, infertility. Recurrent varicocele, which develops after previous surgical or minimally invasive interventions, poses additional challenges for patients, negatively affecting their quality of life and requiring new therapeutic approaches. Therefore, studying recurrent varicocele, understanding its pathogenetic mechanisms, and developing effective treatment strategies is both scientifically and clinically important. Modern medicine emphasizes the preservation of testicular function and the restoration of reproductive potential, as well as improving patients' psychological well-being, making pathogenetically guided approaches to recurrent varicocele highly relevant.

Objective

The main objective of this article is to analyze the mechanisms and pathogenetic basis of recurrent varicocele development, assess its clinical consequences, and review contemporary surgical, endovascular, and conservative treatment methods. Additionally, the study aims to identify strategies for preventing recurrence and preserving patients' reproductive health, providing evidence-based recommendations for clinical practice.

Main part

Recurrent varicocele is a widespread urological condition affecting a significant proportion of men, especially those in reproductive age. Epidemiological studies suggest that varicocele occurs in approximately 15–20% of the male population, while recurrence rates after surgery range between 5% and 15% depending on the technique employed. Clinically, recurrent varicocele can present with testicular pain, heaviness, or atrophy, as well as impaired semen parameters, including oligospermia and asthenospermia. The recurrence often imposes a psychological burden on patients due to concerns about fertility and overall reproductive potential. Early detection of recurrence is challenging, as mild cases may be asymptomatic but still negatively impact spermatogenesis. Ultrasonography with Doppler evaluation is considered the gold standard for diagnosis, allowing for the assessment of venous reflux and testicular volume. Physical examination alone may underestimate the prevalence of recurrence, particularly in cases with subclinical varicoceles. Identifying high-risk groups, such as patients with complex venous anatomy or those with prior inadequate surgical correction, is crucial for prevention and timely management. Understanding the epidemiological and clinical aspects forms the basis for designing effective pathogenetically driven treatment strategies. Comprehensive knowledge of recurrence patterns enables clinicians to tailor therapeutic interventions to individual patient profiles. Moreover, understanding epidemiology aids in resource allocation for urological services and reproductive health programs. Studies indicate that bilateral recurrence may occur, though unilateral left-sided varicocele remains most common. Long-term follow-up is essential to detect late recurrences, which may appear years after initial surgery. Overall, epidemiological insight combined with clinical evaluation provides the foundation for evidence-based management of recurrent varicocele. Effective patient counseling relies on an understanding of the natural history and potential complications of untreated recurrence. Proper documentation of clinical outcomes helps refine surgical techniques



and improve success rates. Hence, studying the epidemiology and clinical significance of recurrent varicocele is fundamental to advancing both research and clinical practice.

Recurrent varicocele develops as a result of complex pathophysiological processes involving venous valve insufficiency, collateral venous pathways, and abnormal venous anatomy. Venous stasis leads to increased hydrostatic pressure in the pampiniform plexus, contributing to testicular hypoxia and oxidative stress. Hypoxia triggers the generation of reactive oxygen species, which can damage sperm DNA and impair spermatogenesis. In addition, impaired venous drainage results in elevated scrotal temperature, further compromising testicular function. Pathogenetically, chronic venous congestion may lead to interstitial fibrosis and Leydig cell dysfunction, affecting testosterone synthesis. Anatomical variations, such as multiple spermatic veins or retroperitoneal collaterals, increase the risk of recurrence if inadequately addressed during initial surgery. Hormonal alterations, including elevated gonadotropins and reduced intratesticular testosterone, have been reported in recurrent cases. Microcirculatory changes in the testicular parenchyma exacerbate oxidative damage and impair nutrient delivery. Endothelial dysfunction within the testicular veins may contribute to persistent reflux and recurrence. Understanding these mechanisms is critical for designing targeted interventions aimed at restoring normal venous flow. Experimental studies have demonstrated that oxidative stress markers correlate with the severity of venous reflux and testicular damage. Preoperative identification of high-risk anatomical features through imaging can reduce recurrence rates. Pathophysiological insights also inform postoperative monitoring and early intervention strategies. Novel pharmacological approaches aimed at reducing oxidative stress are being investigated as adjuncts to surgical management. Overall, a comprehensive understanding of venous stasis and its systemic consequences underpins the pathogenetic rationale for treatment. By addressing both mechanical and biochemical factors, clinicians can improve long-term outcomes for patients with recurrent varicocele. The interplay between venous anatomy, reflux dynamics, and testicular physiology forms the core of modern pathogenetic management strategies. Recognizing these processes emphasizes the importance of precision in both surgical technique and postoperative care.

Accurate diagnosis of recurrent varicocele is essential for effective management and relies on a combination of clinical examination and advanced imaging modalities. Physical examination remains the first-line assessment, including palpation in standing and supine positions to detect dilated veins. However, subclinical recurrences often require color Doppler ultrasonography for confirmation. Doppler studies evaluate the presence of venous reflux, peak retrograde flow, and vein diameter, providing quantitative data for clinical decision-making. Grading systems, such as the Dubin and Amelar classification, are commonly employed to standardize clinical and imaging findings. Recurrent varicocele can be classified as subclinical, clinical grade I–III, or bilateral based on anatomical and hemodynamic parameters. Scrotal thermography and venography may serve as adjunctive tools in complex or ambiguous cases. Non-invasive methods allow monitoring of postoperative patients to detect early recurrence before clinical symptoms manifest. MRI and CT venography are reserved for atypical anatomy or multiple recurrences where standard imaging is insufficient. Understanding the type and severity of recurrence guides the selection of therapeutic approach, whether surgical, endovascular, or combined. Accurate diagnostic assessment also helps predict the potential improvement in semen parameters and testicular function post-treatment. Standardized protocols reduce interobserver variability and improve outcome comparisons across studies. Timely diagnosis minimizes progression of testicular damage and preserves fertility potential. Ultimately,



a systematic diagnostic approach combining clinical, imaging, and functional assessment forms the foundation for evidence-based management of recurrent varicocele. Continuous refinement of imaging techniques enhances early detection, guides individualized therapy, and contributes to improved long-term outcomes.

Surgical intervention remains the primary treatment for recurrent varicocele, aiming to ligate or embolize incompetent veins while preserving testicular arterial supply and lymphatic drainage. Open inguinal, subinguinal, and retroperitoneal approaches are commonly employed, each with specific advantages and limitations. Microsurgical subinguinal varicocelectomy has demonstrated lower recurrence rates due to superior visualization of veins and preservation of arteries. Laparoscopic and robotic-assisted techniques offer minimally invasive alternatives with reduced postoperative pain and faster recovery. Embolization and sclerotherapy serve as effective endovascular options, particularly for patients with prior multiple surgeries. The choice of technique depends on anatomical considerations, prior interventions, and surgeon expertise. Clinical studies indicate that recurrence rates vary according to technique, with microsurgical methods achieving the lowest rates. Complications, such as hydrocele formation, testicular atrophy, or nerve injury, are minimized by meticulous surgical technique and intraoperative identification of structures. Postoperative monitoring with Doppler ultrasonography ensures early detection of residual or recurrent reflux. The timing of surgery is crucial, particularly in adolescents, to prevent long-term testicular damage. Patient selection and preoperative counseling enhance adherence and satisfaction with treatment outcomes. Comparative studies highlight that combination approaches, integrating microsurgery with endovascular embolization, may be beneficial in complex recurrences. Adequate hemostasis and careful ligation of collateral veins reduce recurrence risk. Surgical success is measured not only by anatomical correction but also by improvement in semen parameters and fertility potential. Training and experience of the surgical team are critical for minimizing technical failures. Evidence-based guidelines recommend individualized treatment planning based on the patient's age, fertility goals, and prior surgical history. Long-term follow-up is essential to assess functional outcomes and prevent late recurrence. Ultimately, surgical management remains the cornerstone of recurrent varicocele treatment, guided by pathophysiological understanding and tailored to patient-specific factors.

Endovascular techniques, including percutaneous embolization and sclerotherapy, offer minimally invasive alternatives for recurrent varicocele, especially in patients with complex anatomy or prior surgery. These procedures involve catheterization of the internal spermatic vein and injection of embolic agents to occlude incompetent veins, thereby preventing reflux. Embolization can be performed using coils, sclerosants, or a combination, with procedural success rates ranging from 85% to 95%. Minimal invasiveness reduces postoperative pain, shortens recovery time, and minimizes scarring compared to open surgery. Imaging guidance, such as fluoroscopy or digital subtraction angiography, is essential for precise navigation and effective vein occlusion. Endovascular approaches are particularly advantageous in cases of multiple collaterals or inaccessible veins due to previous interventions. Recurrence rates after embolization are comparable to microsurgical techniques in selected patients. Complications are generally minor, including local hematoma, coil migration, or transient pain. Patient selection is critical, as anatomical variations or occluded veins may limit access. Follow-up Doppler ultrasonography evaluates procedural success and ensures the absence of persistent reflux. Endovascular procedures can be repeated if necessary without significant additional risk. They also allow bilateral treatment in a single session, which is beneficial in patients with bilateral



recurrence. Combining endovascular techniques with microsurgical ligation may optimize outcomes in challenging cases. Pre-procedural counseling includes discussion of fertility outcomes and potential complications. Training and experience of interventional radiologists impact technical success. Advances in embolic materials and imaging modalities continue to improve safety and efficacy. Overall, endovascular therapy represents an essential component of modern, pathogenetically guided management of recurrent varicocele. Integration with conventional surgical strategies enhances treatment flexibility and patient-specific customization. Evidence supports its inclusion in comprehensive care plans for men with recurrent varicocele, particularly when conventional surgery is challenging or contraindicated.

Although surgical and endovascular interventions remain the mainstay of treatment, conservative and pharmacological approaches play an adjunctive role in recurrent varicocele management. Pharmacological strategies aim to mitigate oxidative stress, improve microcirculation, and support testicular function. Antioxidants, such as vitamin C, vitamin E, and coenzyme Q10, have been shown to reduce reactive oxygen species and improve sperm quality. Anti-inflammatory agents may help alleviate testicular discomfort associated with venous congestion. Conservative measures, including scrotal support, avoidance of prolonged standing, and lifestyle modifications, may reduce symptom severity. Pharmacological therapy is typically considered in patients with mild recurrence or those who are not surgical candidates. Evidence suggests that combination therapy with antioxidants and surgical or endovascular intervention may enhance outcomes. Regular monitoring of semen parameters and hormonal profiles guides therapy efficacy. Patient education regarding the progressive nature of recurrent varicocele is essential for adherence to conservative measures. Nutritional support and management of comorbidities, such as obesity or metabolic syndrome, can indirectly improve venous health. Emerging pharmacological agents targeting endothelial function and venous tone are under investigation. Conservative management alone rarely resolves anatomical recurrence but can stabilize testicular function and delay progression. Personalized therapy plans based on severity, fertility goals, and comorbid conditions optimize patient outcomes. Understanding the pathogenetic basis of varicocele recurrence allows for targeted adjunctive therapy. Integration of conservative approaches into multidisciplinary care supports overall reproductive health. Clinical guidelines recommend adjunctive pharmacotherapy as part of comprehensive, evidence-based management. Long-term follow-up ensures timely escalation to surgical or endovascular options if required. Ultimately, conservative and pharmacological measures complement invasive interventions, aiming to preserve testicular function and enhance quality of life for patients with recurrent varicocele.

Preventing recurrent varicocele is critical to maintaining long-term testicular function and reproductive potential. Surgical precision, including meticulous identification and ligation of all incompetent veins, is paramount in reducing recurrence. Preoperative imaging allows mapping of collateral veins and assessment of anatomical variations to guide operative planning. Microsurgical techniques, with magnification and careful preservation of arteries and lymphatics, significantly reduce recurrence rates and postoperative complications. Endovascular strategies can be employed prophylactically in high-risk patients or complex anatomies. Standardized postoperative protocols, including early Doppler ultrasonography, enable detection of residual reflux and timely intervention. Patient education on activity restrictions and scrotal support enhances recovery and reduces recurrence risk. Long-term follow-up is essential, as late recurrences may occur years after initial intervention. Documentation of outcomes informs refinements in surgical technique and helps establish best practice guidelines. Multidisciplinary



collaboration among urologists, interventional radiologists, and reproductive specialists ensures comprehensive care. Identification of high-risk groups, such as adolescents or patients with bilateral varicocele, guides preventive strategies. Lifestyle modifications, including weight management and avoidance of prolonged standing, may contribute to venous health. Regular assessment of semen parameters and hormonal function postoperatively evaluates treatment efficacy. Early identification and management of complications, such as hydrocele or testicular atrophy, prevent long-term morbidity. Evidence-based prophylactic strategies reduce the likelihood of reoperation and improve patient satisfaction. Incorporation of emerging imaging technologies enhances anatomical visualization and procedural planning. Overall, prevention and vigilant monitoring form integral components of pathogenetically guided recurrent varicocele management. Optimizing operative technique, follow-up, and patient education ensures durable outcomes. Preventive approaches not only preserve fertility but also minimize psychological and financial burden on patients. Integrating these strategies into standard care protocols enhances long-term reproductive and testicular health outcomes.

Future perspectives in recurrent varicocele management focus on refining pathogenetically based interventions and enhancing fertility outcomes. Advances in microsurgical and endovascular technologies continue to reduce recurrence rates and postoperative complications. Research into the molecular and cellular mechanisms underlying venous incompetence and oxidative stress may reveal novel pharmacological targets. Biomarkers for early detection of testicular hypoxia and DNA damage could guide timely intervention and improve prognosis. Integration of imaging modalities, such as 3D venography and Doppler thermography, enables precise anatomical mapping and individualized treatment planning. Artificial intelligence and machine learning approaches may predict recurrence risk and optimize surgical strategy selection. Longitudinal studies evaluating reproductive and hormonal outcomes provide evidence for best practice recommendations. Patient-centered research emphasizes quality of life, psychological well-being, and fertility preservation. Novel minimally invasive techniques, including robot-assisted surgery and targeted embolization, are under investigation for complex recurrences. Combination therapies, integrating surgical, endovascular, and pharmacological approaches, offer potential for improved outcomes. International collaboration and multicenter trials can standardize treatment protocols and enhance evidence-based practice. Education and training programs for urologists and interventional radiologists support skill development and reduce technical failures. Genomic and proteomic studies may identify individuals at high risk of recurrence. Personalized medicine approaches hold promise for optimizing treatment efficacy while minimizing adverse effects. Postoperative rehabilitation and lifestyle guidance remain essential components of comprehensive care. Ethical considerations in fertility preservation guide research and clinical decision-making. Public health initiatives may improve awareness, early detection, and intervention in at-risk populations. Ultimately, continued research and innovation will advance pathogenetically grounded management of recurrent varicocele. Emphasis on evidence-based, individualized strategies ensures improved reproductive health outcomes for affected men.

Conclusion

Recurrent varicocele represents a significant clinical challenge due to its impact on testicular function, sperm quality, and male fertility. The pathogenesis of recurrence involves venous valve insufficiency, collateral vein formation, venous stasis, hypoxia, and oxidative stress, which together compromise spermatogenesis and endocrine function. Accurate diagnosis using a



combination of physical examination, Doppler ultrasonography, and advanced imaging modalities is essential for appropriate treatment planning. Management strategies should be guided by pathogenetic principles, incorporating surgical, endovascular, and conservative approaches tailored to individual patient anatomy, prior interventions, and fertility goals. Microsurgical varicocelectomy remains the gold standard for minimizing recurrence, while endovascular techniques provide minimally invasive alternatives, particularly in complex cases or after prior surgery. Adjunctive conservative and pharmacological therapies, including antioxidants and lifestyle modifications, support testicular function and reduce symptom severity. Prevention and vigilant postoperative monitoring are crucial for early detection of residual reflux and reducing the likelihood of further recurrence. Future perspectives emphasize the integration of molecular insights, advanced imaging, and personalized medicine approaches to optimize outcomes and preserve reproductive potential. Overall, a comprehensive, pathogenetically grounded approach ensures effective management of recurrent varicocele, enhances fertility outcomes, and improves the quality of life for affected men.

References

1. Akhmadov, S. R., & Tursunov, B. A. (2020). Pathophysiological basis of varicocele and its recurrence. *Uzbekistan Journal of Urology*, 15(2), 45–52.
2. Abdullayev, M. S., & Karimov, D. F. (2019). Male reproductive health and varicocele. *Medicine and Health*, 8(4), 12–20.
3. Marmar, J. L. The pathophysiology of varicocele and its effect on male fertility. *Human Reproduction Update*, 7(5), 452–462.
4. Shukurov, B. A. (2018). Surgical treatment of varicocele and its efficacy. *Scientific Journal of Uzbekistan Medical Academy*, 10(3), 33–40.
5. Agarwal, A., Deepinder, F., Cocuzza, M., Short, R. A., & Evenson, D. P. (2009). Effect of varicocele repair on sperm DNA integrity in subfertile men. *Fertility and Sterility*, 91(5), 1787–1794.
6. Karimova, S. E., & Rakhmonov, T. M. (2021). Prevention and management of recurrent varicocele. *Tashkent Medical Journal*, 12(1), 25–33.
7. Baazeem, A., & Belker, A. (2014). Varicocele: Contemporary surgical techniques and outcomes. *Arab Journal of Urology*, 12(1), 12–21.
8. Chen, S., & Li, Y. (2020). Endovascular treatment of recurrent varicocele: Techniques and results. *Journal of Vascular and Interventional Radiology*, 31(2), 213–220.
9. Ramasamy, R., & Schlegel, P. N. (2008). Microsurgical treatment of recurrent varicocele. *Asian Journal of Andrology*, 10(1), 107–112.