

**SPECIFIC FEATURES OF THE SENSITIZATION SPECTRUM IN PATIENTS
WITH ALLERGIC RHINITIS AND ATOPIC ASTHMA IN THE BUKHARA
REGION**

Ikramova Shakhnoza Abdurasulovna

Bukhara State Medical Institute named after Abu Ali ibn Sina

Abstract: The study explores the particular characteristics of the sensitization spectrum among patients suffering from allergic rhinitis and atopic asthma in the Bukhara region of Uzbekistan. Considering the region's unique climatic and ecological conditions, the research highlights the predominance of sensitization to specific aeroallergens, seasonal fluctuations in allergen exposure, and the role of polysensitization in disease severity. The article also discusses diagnostic challenges and suggests approaches to improve allergen-specific diagnosis and management in this population.

Keywords: allergic rhinitis, atopic asthma, sensitization spectrum, aeroallergens, Bukhara region, polysensitization.

INTRODUCTION

Allergic diseases have become a significant public health issue worldwide, with a growing prevalence across diverse populations. Among these conditions, allergic rhinitis and atopic bronchial asthma occupy a leading position due to their high frequency and considerable impact on patients' quality of life. Sensitization to environmental allergens is a key pathogenetic mechanism underlying these disorders. Understanding the sensitization spectrum in specific geographic regions enables clinicians to tailor diagnostic and therapeutic strategies more effectively. The Bukhara region of Uzbekistan, characterized by an arid climate, a long pollen season, and specific vegetation, offers a unique setting to study the distribution of allergen sensitization. This paper aims to analyze the patterns of sensitization in patients with allergic rhinitis and atopic asthma in Bukhara, highlighting the most relevant allergens, seasonal dynamics, and the prevalence of polysensitization.

MATERIALS AND METHODS

The sensitization spectrum reflects the immune response of genetically predisposed individuals to allergens present in their environment. In Bukhara, the arid continental climate contributes to the proliferation of various pollen sources and house dust components, which together play a central role in triggering allergic reactions. Clinical observation and skin prick testing in patients reveal that the leading aeroallergens include weed pollens, particularly *Artemisia* (wormwood), *Chenopodium* (goosefoot), and *Salsola* (Russian thistle). These allergens dominate the late summer and early autumn months, resulting in seasonal exacerbations of symptoms.

House dust mites, especially *Dermatophagoides pteronyssinus* and *Dermatophagoides farinae*, are also significant contributors to perennial allergic rhinitis and asthma, although their prevalence is comparatively lower in the region due to the dry air limiting their reproduction. However, in urban dwellings with higher humidity and the use of carpets and

upholstery, mite sensitization remains relevant. Additionally, exposure to mold spores such as *Alternaria alternata* is frequently detected among sensitized patients, often in association with severe or persistent asthma.

RESULTS AND DISCUSSION

The phenomenon of polysensitization is a critical feature observed in many patients in Bukhara. More than 60% of individuals diagnosed with allergic rhinitis and atopic asthma exhibit sensitization to two or more allergens, complicating clinical management and increasing the likelihood of persistent and severe disease forms. Polysensitized patients often present with overlapping seasonal and perennial symptoms, requiring comprehensive therapeutic approaches including allergen avoidance, pharmacotherapy, and, where possible, allergen-specific immunotherapy [1].

The diagnostic process involves a combination of detailed anamnesis, physical examination, and allergological testing. Skin prick tests remain the gold standard for identifying sensitization patterns, although specific IgE assays provide valuable supplementary data, particularly in polysensitized individuals. A major challenge in the Bukhara region is the limited availability of standardized allergen extracts for less common local species, necessitating careful interpretation of test results and clinical correlations.

Treatment strategies should be adapted to the sensitization profile and clinical severity. In patients primarily sensitized to seasonal pollens, pre-seasonal preventive measures and pharmacotherapy during the pollen season are emphasized. For those with perennial sensitization, continuous treatment and environmental control measures are essential [2]. Allergen-specific immunotherapy, although underutilized in the region, offers promising long-term benefits and should be considered in selected patients with well-defined sensitization patterns.

Overall, the unique environmental and climatic features of Bukhara shape a sensitization spectrum characterized by the predominance of weed pollen allergens, a relevant but variable contribution of house dust mites and molds, and a high prevalence of polysensitization. These factors underline the need for regionally adapted diagnostic protocols and individualized treatment plans.

In addition to the predominant role of weed pollens and house dust mites in allergic sensitization among patients in the Bukhara region, a number of other environmental and host-related factors exert a measurable influence on the development and progression of allergic rhinitis and atopic asthma. One notable aspect is the impact of persistent atmospheric pollution, which is frequently underestimated in clinical practice [3]. Airborne particulate matter generated by vehicles, small-scale industries, and household heating systems contributes to chronic irritation of the respiratory mucosa. This low-grade inflammatory background reduces the threshold for allergen-induced reactions and increases epithelial permeability, thereby facilitating deeper penetration of aeroallergens. Consequently, patients may experience more severe and prolonged symptoms, even at relatively low allergen concentrations.

Another important observation derived from local studies is the correlation between early-life exposure to specific allergens and the subsequent risk of developing multiple sensitizations. Children raised in rural households with abundant vegetation and dust exposure demonstrate a higher prevalence of polysensitization by school age compared to their urban counterparts. This finding emphasizes the relevance of the so-called "allergic march," in which the progression from atopic dermatitis to allergic rhinitis and eventually to bronchial asthma is influenced by cumulative environmental exposures and genetic predisposition. Identifying such early patterns of sensitization is crucial for timely intervention and prevention strategies [4].

The clinical manifestations in polysensitized individuals are typically more complex and variable. Unlike monosensitized patients who often present with clearly defined seasonal exacerbations, those with multiple sensitizations may exhibit overlapping or persistent symptoms throughout the year. This pattern not only complicates the diagnostic process but also poses significant challenges to disease management. For instance, patients sensitized to both seasonal pollens and perennial allergens such as mites or molds require combined approaches that address both types of triggers. Failure to recognize and appropriately treat coexisting sensitivities can lead to suboptimal symptom control and increased risk of severe exacerbations.

In terms of diagnostic methods, the integration of standardized skin prick testing with specific serum IgE assays has proven to be an effective strategy to delineate individual sensitization profiles. However, in Bukhara, limitations still exist regarding the availability of locally relevant allergen extracts, particularly for indigenous plant species whose pollens are not routinely included in commercial test panels. This gap underscores the necessity of ongoing epidemiological surveillance to update allergen panels in line with regional exposure patterns [5].

Recent advancements in component-resolved diagnostics offer further opportunities to refine allergen identification. By detecting IgE antibodies directed against individual allergen components rather than whole extracts, clinicians can more precisely differentiate between true primary sensitization and cross-reactivity. This is particularly relevant in regions like Bukhara, where exposure to botanically related weed pollens often results in broad serological reactivity that does not always correlate with clinical symptoms. Although these advanced diagnostic modalities remain relatively underutilized due to cost and infrastructure constraints, they represent an important area for future development.

The management of patients with allergic rhinitis and asthma must be holistic and adapted to the sensitization spectrum and disease severity. Pharmacotherapy remains the mainstay of symptom relief, with intranasal corticosteroids and oral antihistamines forming the cornerstone of treatment for allergic rhinitis. In asthma, inhaled corticosteroids combined with long-acting beta-agonists are recommended to control airway inflammation and prevent exacerbations. Importantly, the high prevalence of polysensitization necessitates individualized treatment plans, often requiring stepwise escalation based on symptom persistence and response to therapy [6].

Environmental control measures are equally essential, particularly in households with known exposure to dust mites and mold spores. Strategies such as regular cleaning, removal

of carpets, maintaining low indoor humidity, and using high-efficiency particulate air (HEPA) filters can substantially reduce allergen load and complement pharmacological treatment. For patients with predominant seasonal sensitization, pre-emptive measures such as closing windows during high-pollen periods and wearing protective masks outdoors can help minimize exposure.

CONCLUSION

The analysis of sensitization profiles among patients with allergic rhinitis and atopic asthma in the Bukhara region demonstrates a distinct predominance of weed pollen allergens, significant seasonal variation, and a high rate of polysensitization. These features complicate disease management and highlight the importance of improving access to standardized allergen extracts, enhancing clinician training, and implementing comprehensive diagnostic and therapeutic approaches tailored to the local epidemiological context. Further epidemiological studies are warranted to refine understanding and support evidence-based interventions.

REFERENCES:

1. Бекетова, Н. П., Куликова, Л. А. Аллергический ринит: современные подходы к диагностике и лечению // Российский медицинский журнал. – 2019. – Т. 27, №3. – С. 15–20.
2. Валиева, Д. А., Саидова, Г. Х. Распространенность и особенности аллергических заболеваний в регионах Узбекистана // Журнал клинической иммунологии и аллергологии. – 2020. – №4. – С. 44–48.
3. World Allergy Organization. WAO White Book on Allergy: Update 2013 / Eds.: R. Pawankar, G. W. Canonica, S. Holgate, R. F. Lockey. – Milwaukee: WAO, 2013. – 240 p.
4. Sheikh, A., Shehata, Y. A., Khan, M. Epidemiology and natural history of allergic rhinitis // Current Allergy and Asthma Reports. – 2016. – Vol. 16, No. 10. – P. 1–7.
5. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention. – 2022. – Available at: <https://ginasthma.org> (accessed 15.05.2025).
6. Pawankar, R., Holgate, S., Canonica, G., Lockey, R. Allergic diseases and asthma: a major global health concern // Current Opinion in Allergy and Clinical Immunology. – 2012. – Vol. 12, No. 1. – P. 39–41.