

STOMATOLOGY OF CHILDREN WITH ATOPIC DERMATITIS STATUS

Murtazaev S.S., Eshkulova Sh.B.

Tashkent State Dental Institute

eshkulovashaxzoda@gmail.com

Abstract. In recent years, allergic diseases have been on the rise in the world. Atopic dermatitis takes the leading place among these diseases. Atopic dermatitis causes the development of bronchial asthma, pollinosis, urticaria, Quincke's tumor, allergic rhinitis in 70% of cases.

Key words: Dermatitis, allergic diseases, dental diseases, diagnosis of dermatitis

Young children have a high risk of developing dental diseases due to poor hygiene and poor physical condition.[2]

When planning prevention, it is necessary to take into account the high risk of occurrence and development of dental diseases in children [1]. Children with allergopathology are especially at risk.[5,6]

According to the literature, it is present in patients with allergopathology

Various pathological changes in the dental-chewing system are: swelling of the mucous membrane of the oral cavity, aphthous stomatitis, desquamative glossitis, exfoliative and eczematous cheilitis. changes in the composition of oral fluid and gum fluid, oral cavity immunity, systemic processes, hemodynamic disorders in periodontal tissues are detected. [3,4] Including a large number of children with atopic dermatitis. it is necessary to emphasize that and the disease has a relapsing nature. [9,10] Such children receive high-dose drug therapy for several years, which in turn causes local and systemic negative effects on the tissues of the oral cavity. can have an effect.[1]

It is noted that general somatic diseases are one of the factors that keep the frequency of morphological and functional disorders in the teeth and jaw system in children [2]. It has been determined by scientists that the functional state of the tooth-jaw system and other systems is connected through inter-system integration. [1, 7].

The purpose of the study. Study of dental status in children with atopic dermatitis.

Material and methods. For research, 25 children of different ages with a diagnosis of atopic dermatitis, registered in the Republican center of specialized allergology and clinical immunology, were examined (by gender, 15 girls - 60%, 10 boys child - 40%).

According to life anamnesis indicators, 60% of children have severe allergic anamnesis.

During the study, the patients underwent a clinical examination of the oral cavity. Also, the condition of the teeth, mucous membrane of the oral cavity, pricus, and vulva were

studied. Teeth were evaluated according to caries intensity, KPO indices. The level of oral hygiene was studied with the Fedorova-Volodkina hygiene index. The condition of periodontal tissues was assessed by PMA and PI indices with a special button-shaped probe.

Results and discussions. According to the results of the study, the prevalence of caries in 1.5-6-year-old children diagnosed with atopic dermatitis was 55%, 7-11 years old - 80%, 12-15 years old - 100%. Caries intensity 1.5-6 years-2.4;

7-11 years-4.4; 12-15 years old - showed 2.9.

According to our results, the factor that causes high caries prevalence and caries intensity in children with atopic dermatitis is the effect of the disease on the resistance of dental hard tissues. It is also worth noting that 12-15-year-old children have a lower level of caries intensity, because this physiological baby teeth have changed, but the level of caries prevalence is high. [3].


When determining the position of the lumps in the framework of teeth-jaw anomalies, in 37% of the examined patients with milk and mixed pricus, the lumps are located below. The percentages of those with permanent pricus are 8% with a short wedge-shaped distal occlusion, 4% with a mesial occlusion after frenuloplasty, and 4% with a deep occlusion. In the remaining 47% of children, no pricus anomalies were detected. Pricus position, dentition position, and occlusal position were age-appropriate.

When assessing the oral hygiene condition, the following indicators were found: 40% of children had very poor hygiene, 12% - poor, 20% - unsatisfactory, 8% - satisfactory, and 20% - good.

When periodontal tissue status was determined by PI, 26% had mild periodontal disease, 48% were at risk of developing, and the remaining 26% had no disease.

When the level of gingivitis of the children in the study was assessed according to the PMA index, 12% had a severe type, 20% had a moderate level, and 68% had no symptoms.

In 64% of children, various forms of cheilitis, especially exfoliative and atopic cheilitis, changes in the corner of the mouth, and dryness of the lips and tongue were found in the red border of the lips. This result was 1.5 times higher compared to a similar study [11]. In 16% of children, the tongue is covered with scurvy.



Conclusion.

1. According to the results of the research, children with atopic dermatitis are more likely to get dental diseases due to the fact that they received general medical treatment, especially oral drugs. Therefore, it is necessary to develop preventive measures for dental diseases in such patients.

2. In order to reduce the risk of dental diseases in children with allergic diseases, especially atopic dermatitis, it is necessary to control the observance of hygiene rules.

References:

1. Адмакин О.И. Стоматологический статус детей 12-летнего возраста, страдающих аллергической патологией // Стоматология детского возраста. – 2007. – № 2.
2. Овсянникова О.Б. Атопические дерматиты в амбулаторной практике // Медицинский совет. – 2012. – № 3.
3. Попруженко Т.В., Терехова Т.Н. Профилактика основных стоматологических заболеваний. – М.: МЕДпресс – инфо, 2009.
4. Аллергология и иммунология: клинические рекомендации для педиатров / А.А. Баранов [и др.]. – М.: Союз педиатров России. – 3-е изд., испр. и доп., 2011.
5. Узиков, О.Ж. Атопический дерматит: новые аспекты этиопатогенеза, клиники, диагностики, терапии и профилактики: учебное пособие / О.Ж. Узиков, Ж.К. Муратова, Б.Д. Кудаяров. – Ош, 2012.
6. Балаболкин, И.И. Актуальные проблемы аллергологии детского возраста на современном этапе / И.И. Балаболкин // Педиатрия. – 2012. – Т. 91, № 3.
7. Боронбаева, Э.К. Современные методологические аспекты эпидемиологических и клинических исследований аллергической патологии у детей: автореф. дисс. ... канд. мед. наук. – Бишкек, 2005.

8. Кожевникова, С.Л. Клинико-эпидемиологические особенности бронхиальной астмы и аллергического ринита у школьников г. Чебоксары: автореф. дис. ... канд. мед. наук. – Казань, 2001.
9. Лысикова, И.В. Распространенность аллергических заболеваний у детей по результатам мультицентровых исследований в рамках международной программы ISAAC: автореф. дис. канд. мед. наук. – М., 1999
10. Луткова, Т.С. Клинико-эпидемиологические аспекты атопического дерматита у школьников города Чебоксары: автореф. дис. ... канд. мед. наук. – Чебоксары, 2005.
11. Особенности стоматологического статуса и анализ микрофлоры ротовой полости у детей с аллергической патологией. Современная медицина:актуальные вопросы, No2-3, 2016